

OTC Medicine - Quiz 3

LaRue Medical Literacy Exercises

Drug Facts	
Active Ingredient (in each tablet) <i>Chlorpheniramine Maleate 2 mg</i>	Purpose <i>Antihistamine</i>
Uses temporarily relieves these symptoms due to allergies <div style="display: flex; justify-content: space-around; margin-top: 5px;"> ■ <i>sneezing</i> ■ <i>runny nose</i> </div>	
Warnings Ask a doctor before use if you have <div style="display: flex; justify-content: space-around; margin-top: 5px;"> ■ <i>glaucoma</i> ■ <i>a breathing problem such as emphysema</i> </div>	
When using this product <div style="margin-top: 5px;"> ■ <i>drowsiness may occur</i> ■ <i>be careful when driving a motor vehicle or operating machinery</i> </div>	
Directions	
Adults and children 12 and over	<i>take 2 tablets every 4 to 6 hours not more than 12 tablets in 24 hours</i>
Children 6 years to under 12 years	<i>take 1 tablet every 4 to 6 hours not more than 6 tablets in 24 hour</i>
Children under 6 years	<i>ask a doctor</i>

1. What symptom does this relieve?
 - A. antihistamine
 - B. glaucoma
 - C. sneezing
 - D. drowsiness

2. How much medicine should an adult take?
 - A. one tablet
 - B. two tablets
 - C. ask a doctor
 - D. not more than 6 tablets

3. Who should ask a doctor before using this medicine?
 - A. someone with allergies
 - B. someone with drowsiness
 - C. someone with a runny nose
 - D. someone with glaucoma

4. What is a side effect of this medicine?
 - A. sneezing
 - B. glaucoma
 - C. allergies
 - D. drowsiness

5. How much medicine should a 9-year-old take in one day?
 - A. not more than 12 tablets
 - B. not more than 6 tablets
 - C. ask a doctor
 - D. 1 tablet

Drug Facts	
Active Ingredient (in each tablet)	Purpose
Pseudoephedrine HCL 30 mg.....	Nasal Decongestant
Uses temporarily relieves nasal congestion due to	
<input type="checkbox"/> common cold <input type="checkbox"/> respiratory allergies	
Warnings	
Ask a doctor before use if you have	
<input type="checkbox"/> high blood pressure <input type="checkbox"/> heart disease <input type="checkbox"/> thyroid disease <input type="checkbox"/> diabetes	
When using this product	
<input type="checkbox"/> do not use more than directed <input type="checkbox"/> Stop use and ask a doctor if you get nervous, dizzy or sleepless	
Directions	
Adults and children 12 and over	take 2 tablets every 4 to 6 hours not more than 4 doses in 24 hours
Children 6 years to under 12 years	take 1 tablet every 4 to 6 hours not more than 4 doses in 24 hours
Children under 6 years	ask a doctor

6. What symptom does this medicine treat?
- A. high blood pressure
B. nasal congestion
C. nervousness
D. diabetes
7. How much medicine should a child under 6 take?
- A. two tablets
B. one tablet
C. ask a doctor
D. 4 doses
8. Who should ask a doctor before using this?
- A. someone with nasal congestion
B. someone with allergies
C. someone with a common cold
D. someone with thyroid disease
9. How much medicine can an adult take in 24 hours?
- A. 2 tablets
B. 8 tablets
C. 4 tablets
D. ask a doctor
10. What is a side effect of this medicine?
- A. dizziness
B. respiratory allergies
C. diabetes
D. nasal congestion